

# 表4 タミフル投与時期と胎児・新生児異常との関連

	流産	死産	早産	低出生体重児 (2500g未満児)	胎児発育不全 (FGR)	心形態異常	その他の形態 異常	その他の新生 児異常
無影響期	0/5(0%)	—	1/5(20%)	1/5(20%)	0/5	0/5(0%)	0/5	0/5
絶対過敏期	3/59(5.1%)	—	3/59(5.1%)	4/59(6.8%)	0/59	1/59(1.7%)	0/59	0/59
相対過敏期	0	—	4/99(4.0%)	4/99(4.0%)	1/99(1.0%)	1/99(1.0%)	0/99	6/99(6.1%)
比較過敏期	0	—	7/87(8.0%)	10/87(11.5%)	2/87(2.3%)	0/87	1/87(1.1%)	4/87(4.6%)
小計	3/250(0.4%)	—	15/250(6.0%)	19/250(7.6%)	3/250(1.2%)	2/250(0.8%)	1/250(0.4%)	10/250(4.0%)
妊娠中期	—	1/277(0.02%)	22/277(7.9%)	30/277(10.8%)	10/277(3.6%)	7/277(2.5%)	3/277(1.1%)*	12/277(4.3%)
妊娠後期	—	0/250(0%)	9/250(3.6%)	22/250(8.8%)	14/250(5.6%)	5/250(2.0%)	3/250(1.2%)	4/250(1.6%)
総計	3/250(0.4%)	1/627(0.16%)	46/778(5.9%)	71/778(9.1%)	27/778(3.5%)	14/778(1.8%)	7/778(0.9%)	26/778(3.3%)

\*1例は心奇形と重複